arXiv:hep-ph/0607016v2 11 Sep 2006

The rational parts of one-loop QCD amplitudes II: The five-gluon case

Xun Su*[1], Zhi-Guang Xiao†[1,2], Gang Yang‡[1]and Chuan-Jie Zhu§[1,3]

February 2, 2008

[1]Institute of Theoretical Physics, Chinese Academy of Sciences
P. O. Box 2735, Beijing 100080, P. R. China

[2]The Interdisciplinary Center of Theoretical Studies, Chinese
Academy of Sciences, P. O. Box 2735, Beijing 100080, P. R. China

[3]Center of Mathematical Science, Zhejiang University
Hangzhou 310027, P. R. China

Abstract

The rational parts of 5-gluon one-loop amplitudes are computed by using the newly developed method for computing the rational parts directly from Feynman integrals. We found complete agreement with the previously well-known results of Bern, Dixon and Kosower obtained by using the string theory method. Intermediate results for some combinations of Feynman diagrams are presented in order to show the efficiency of the method and the local cancellation between different contributions.

---

*E-mail: ghkcn@itp.ac.cn
†E-mail: zhgxiao@itp.ac.cn
‡E-mail: yangg@itp.ac.cn
§E-mail: zhucj@itp.ac.cn

1 Introduction

In a previous paper [1], we developed a method for computing the rational parts of one-loop amplitudes directly from Feynman integrals. The purpose of this paper is to apply this method to compute the rational parts of 5-gluon one-loop amplitudes. The result agrees with the well-known result of Bern, Dixon and Kosower [2] obtained first by using string-inspired methods. (Other 5-parton amplitudes in massless QCD were later computed by using either standard Feynman diagrammatic technique [3] or supersymmetric decomposition and perturbative unitarity [4].)

The computation of the multi-particle one-loop amplitudes in QCD is a very difficult problem. Even for 4-parton amplitudes the computation is quite non-trivial [5]. For 5-gluon amplitudes a new method was developed [6] by using string theory.

The constant effort to calculate multi-leg one-loop amplitudes lies in the application to the forthcoming experimental program at CERN's Large Hadron Collider (LHC), as there are lots of processes with many particles as final states [7]. We refer the reader to [1] for a discussion and extensive references for the recent efforts in computing the multi-leg one-loop amplitudes and the recent developments inspired by twistor string theory [8, 9].

In order to compute multi-leg one-loop amplitudes in QCD, it is a good strategy [10] to decompose the QCD amplitudes into simpler ones by using the supersymmetric decomposition:

$$A^{QCD} = A^{N=4} - 4A^{N=1 \text{ chiral}} + A^{N=0 \text{ or scalar}}, \tag{1}$$

where A^{QCD} denotes an amplitude with only a gluon circulating in the loop, $A^{N=4,1}$ have the full $N = 4, 1$ multiplets circulating in the loop, and $A^{N=0}$ has only a complex scalar in the loop.

By using the general properties of the one-loop amplitudes, Bern, Dunbar, Dixon and Kosower proved that the supersymmetric amplitudes $A^{N=4,1}$ are completely determined by 4-dimensional unitarity [10], i.e. the amplitudes are completely cut-constructible and the rational part is vanishing (see [10, 1] for more detail explanation). For MHV helicity configurations, explicit results were obtained for $A^{N=4}$ in [10]. The recent development of using MHV vertices to compute one-loop amplitudes leads to many new results for the cut-constructible part [11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 21, 22]. In particular, Bedford, Brandhuber, Spence and Travaglini [11, 15] applied the

MHV vertices to one-loop calculations. Britto, Buchbinder, Cachazo, Feng and Mastrolia [20, 21, 22] developed an efficient technique for evaluating the rational coefficients in an expansion of the one-loop amplitudes in terms of scalar box, triangle and bubble integrals (the cut-constructible part, see [1] for details). By using their technique, it is much easier to calculate the coefficients of box integrals without doing any integration. Recently, Britto, Feng and Mastrolia completed the computation of the cut-constructible terms for all the 6-gluon helicity amplitudes [22].

In order to complete the QCD calculation for the 6-gluon helicity amplitudes, the remaining challenge is to compute the rational parts of the helicity amplitudes with scalars circulating in the loop. In general, we need an efficient and powerful method to compute the rational part of any amplitude.

As we reviewed in [1], there are various approaches [23, 24, 25, 26, 27, 28] to compute the rational part. In particular, Bern, Dixon and Kosower [27, 28] developed the bootstrap recursive approach which has lead to quite general results [29, 30, 31]. In this paper we will use the approach as developed in [1] and apply it to compute the rational parts of the 5-gluon one-loop amplitudes. In another paper [32] we will compute the rational parts of 6-gluon amplitudes in QCD (see also [31]) which are the last missing pieces for the complete partial helicity amplitudes of the 6-gluon one-loop QCD amplitude.

This paper is organized as follows: in Sect. 2, we recall briefly the Feynman diagrams and the Feynman rules, tailored for our computation of the 5-gluon amplitudes. Some simple reduction formulas are recalled briefly in Sect. 3. In Sect. 4 we summarize all the integral formulas we will use in this paper. Then the following 2 sections present the results for the rational parts of the two independent MHV helicity configurations.

2 Notation, the Feynman diagrams and the Feynman rules

A word about notation: we use the same notation as given in [1]. We use $\epsilon_{i(i+1)\cdots(i+n)}$ to denote the composite polarization vectors for sewing trees to the loop.

For the purpose of this paper we consider only the Feynman diagrams and Feynman rules for the one-loop gluon amplitudes with scalars circulating in the loop. We do not follow the usual convention of differentiating different particles by different kinds of lines because there are only two kinds of particles: gluons and scalars, and scalars only appear in the loop.



Figure 1: The Feynman rules for sewing trees to loop. The blob denotes an expansion of the tree amplitude.

For the explicit calculation of one-loop amplitudes by the usual Feynman diagram technique, we can first collect all terms with the same loop structure into one entity. Generally a few cyclicly consecutive external lines are joined in tree diagrams and connected to the same point on the loop (sewing trees to the loop). We denote the sum of all these contributions by $P_{i(i+1)\cdots(i+m-1)}$ for m such external lines. For $m = 1, 2, 3$, the relevant Feynman diagrams are shown in Fig. 1. Explicitly we have:

$$P_i(p) = (\epsilon_i, p) = (\epsilon_i, p - k_i), \tag{2}$$

$$P_{i(i+1)}(p) = (\epsilon_{i(i+1)}, p) - \frac{1}{2}(\epsilon_i, \epsilon_{i+1}), \tag{3}$$

$$P_{i(i+1)(i+2)}(p) = (\epsilon_{i(i+1)(i+2)}, p) - \frac{1}{2}\left((\epsilon_{i(i+1)}, \epsilon_{i+2}) + (\epsilon_i, \epsilon_{(i+1)(i+2)})\right), \tag{4}$$

where $\epsilon_{\ldots}$'s are composite polarization vectors introduced in [1]. The computation of these composite polarization vectors is a simplified version of the general recursive calculation of the tree-level n-gluon amplitudes [33].



Figure 2: All the possible one-loop Feynman diagrams for 5 gluons. The index i runs from 1 to 5 if there is an index i.

Considering all these different tree diagrams just as the same diagram and denoting them by multiple parallel lines attached to the loop, we have only 21 different Feynman diagrams for the 5-gluon one-loop amplitude (with scalars circulating the loop). Some representative diagrams are given in Fig. 2. The counting goes as follows:

- 1 pentagon diagram, the diagram (a);
- 5 box diagrams because there are 5 different ways of combining two consecutive external lines, diagram (b) with $i = 1, \cdots, 5$;
- 10 triangle diagrams which are further divided into 5 two-mass triangle diagrams (diagram (c)) and 5 one-mass triangle diagrams (diagram (d));
- 5 bubble diagrams (e).

It is straightforward to compute the rational part from each diagram for a given helicity configuration. In the next two sections we will review all the formulas needed.

3 Tensor reduction of the one-loop amplitudes

There is a vast literature on this subject [34, 35, 36, 37, 38, 39, 40]. The tensor reduction relations we will use for our calculation of 5-gluon amplitudes are quite simple. It is based on the BDK trick [28] of multiplying and dividing by spinor square roots. For 5-gluon amplitudes it is not so important to make a clever choice of reference momenta. For 6-gluon amplitudes it is important to make a specific choice of the reference momenta to make all tensor reductions simple enough to obtain relatively compact analytic results.



Figure 3: For two adjacent same helicities, the tensor reduction for the combination of two diagrams is even simpler.

For the tensor reduction with only 2 neighboring same helicity external gluons, it is possible to choose the reference momenta to be each other's momenta, i.e. $\epsilon_1 = \lambda_1\tilde{\lambda}_2$, $\epsilon_2 = \lambda_2\tilde{\lambda}_1$.[1] The tensor reduction is done by considering the contributions from 2 diagrams together and the result formula is shown pictorially in Fig. 3. The exact algebraic formula is:

$$\frac{(\epsilon_1, p+k_1)(\epsilon_2, p)}{(p+k_1)^2 p^2 (p-k_2)^2} + \frac{(\epsilon_{12}, p+k_1) - (\epsilon_1, \epsilon_2)/2}{(p+k_1)^2 (p-k_2)^2} = -\frac{1}{p^2} + \frac{1/2}{(p+k_1)^2} + \frac{1/2}{(p-k_2)^2}. \tag{5}$$

For a different choice of reference momenta,

$$\epsilon_4 = \lambda_5\tilde{\lambda}_4, \qquad \epsilon_5 = \eta\tilde{\lambda}_5, \tag{6}$$

---

[1] An overall constant is omitted for the polarization vector which can be easily reinstated at the end of calculation. See [41, 42] for details.



Figure 4: A combination of 2 diagrams with the same adjacent helicity.

the tensor reduction of the 2 diagrams shown in Fig. 4 is given as follows:

$$\begin{aligned} A_a + A_b &= \frac{P_4(p)\, P_5(p-k_4)}{p^2\,(p-k_4)^2\,(p-k_{45})^2} + \frac{P_{45}(p)}{p^2(p-k_{45})^2} \\ &= -\frac{(\eta\tilde{\lambda}_4, p)}{p^2\,(p-k_4)^2} + \frac{\langle \eta\, 5\rangle}{2\,\langle 4\, 5\rangle} \left[\frac{1}{(p-k_{45})^2} - \frac{1}{p^2}\right]. \end{aligned} \tag{7}$$

Setting $\eta = \lambda_4$, the above equation agrees with eq. (5).

4 A summary of the rational parts for triangle and box integrals

For quick reference we list here the explicit formulas for the rational parts of some Feynman integrals which are needed to compute the 5-gluon amplitudes. The derivation can be found in [1].

Firstly, for the bubble integral we have:

$$\begin{aligned} I_2(\epsilon_1, \epsilon_2) &= \int \frac{\mathrm{d}^D p}{i\pi^{D/2}} \frac{(\epsilon_1, p)(\epsilon_2, p)}{p^2(p+K)^2} \\ &= \frac{1}{18}\left((\epsilon_1, K)\,(\epsilon_2, K) - 2\,K^2\,(\epsilon_1, \epsilon_2)\right). \end{aligned} \tag{8}$$

where K is the sum of momenta on one side of the bubble diagram. The above result first appeared in [2].

For 2-mass triangle integral with external momenta $\{k_1, K_2, K_3\}$ and $k_1^2 = 0$, the degree 2 integral is:

$$I_3(\epsilon_1, \epsilon_2) \equiv \int \frac{\mathrm{d}^D p}{i\pi^{D/2}} \frac{(\epsilon_1, p)\,(\epsilon_2, p)}{p^2(p-k_1)^2(p+K_3)^2}$$

$$
\begin{aligned}
&= \frac{1}{2}\,(\epsilon_1,\epsilon_2) + \frac{(K_2^2+K_3^2)}{2(K_2^2-K_3^2)^2}\,(\epsilon_1,k_1)\,(\epsilon_2,k_1) \\
&+ \frac{((\epsilon_1,K_2)\,(\epsilon_2,k_1)-(\epsilon_1,k_1)\,(\epsilon_2,K_3))}{2(K_2^2-K_3^2)}, \qquad (9)
\end{aligned}
$$

$$
I_3(\epsilon_1,\epsilon_2) = \frac{1}{2}\,(\epsilon_1,\epsilon_2) + \frac{(\epsilon_1,K_2)\,(\epsilon_2,k_1)}{2(K_2^2-K_3^2)}, \qquad (\epsilon_1,k_1)=0, \qquad (10)
$$

and the degree 3 integral is:

$$
\begin{aligned}
I_3(\epsilon_i) &\equiv \int \frac{\mathrm{d}^D p}{i\pi^{D/2}}\,\frac{(\epsilon_1,p)\,(\epsilon_2,p-k_1)\,(\epsilon_3,p)}{p^2\,(p-k_1)^2\,(p+K_3)^2} \\
&= \frac{1}{36}\Big((\epsilon_2,4\,K_2-7\,k_1)\,(\epsilon_1,\epsilon_3)-(2\leftrightarrow 3)+4(\epsilon_1,K_2)\,(\epsilon_2,\epsilon_3)\Big) \\
&- \frac{(K_2^2+K_3^2)}{6\,(K_2^2-K_3^2)^2}\,(\epsilon_1,K_2)\,(\epsilon_2,k_1)\,(\epsilon_3,k_1) \\
&- \frac{(\epsilon_1,K_2)\,((\epsilon_2,k_1)\,(\epsilon_3,K_3)-(\epsilon_2,K_2)\,(\epsilon_3,k_1))}{6\,(K_2^2-K_3^2)} \\
&- \frac{(K_2^2+K_3^2)}{12\,(K_2^2-K_3^2)}\,((\epsilon_1,\epsilon_2)\,(\epsilon_3,k_1)+(\epsilon_1,\epsilon_3)\,(\epsilon_2,k_1)), \qquad (11)
\end{aligned}
$$

where ϵ_1 satisfies the physical condition $(\epsilon_1,k_1)=0$ and $\epsilon_{2,3}$ are arbitrary 4-dimensional polarization vectors.

For five point cases we need only 1-mass box integrals. The 1-mass box integrals can be obtained simply by setting the mass of one massive external line to 0 from the 2-mass-easy box integrals. So we list the formulas of two-mass-easy box integrals here. The four external momenta of the 2-mass-easy box diagram are denoted as $\{k_1,K_2,k_3,K_4\}$.

For degree 3 two-mass-easy box integrals, we have:

$$
I_4^{2me}(\lambda_3\tilde{\lambda}_1,\epsilon_2,\epsilon_3) = \frac{\langle 3|K_2|1]}{2}\left[\frac{(\epsilon_2,k_3)\,(\epsilon_3,k_3)}{(K_2^2-t)(K_4^2-s)} - (k_3\to k_1,\, s\leftrightarrow t)\right], \qquad (12)
$$

where $s=(k_1+K_2)^2$ and $t=(K_2+k_3)^2$. If two of the polarization vectors satisfy the physical condition, i.e. $(\epsilon_1,k_1)=0$ and $(\epsilon_3,k_3)=0$, we have

$$
\begin{aligned}
I_4^{2me}(\epsilon_1,\epsilon_2,\epsilon_3) &= -\frac{(\epsilon_1,k_3)\,(\epsilon_3,k_1)}{2\,(k_1,k_3)}\left[\frac{(\epsilon_2,k_3)}{K_2^2-t}+\frac{(\epsilon_2,k_1)}{K_4^2-t}\right] \\
&\quad -\frac{(\epsilon_1,K_2)\,(\epsilon_2,k_1)\,(\epsilon_3,k_1)}{2\,(K_2^2-s)\,(K_4^2-t)} - \frac{(\epsilon_1,k_3)\,(\epsilon_2,k_3)\,(\epsilon_3,K_4)}{2\,(K_2^2-t)\,(K_4^2-s)}. \qquad (13)
\end{aligned}
$$

In order to give the formulas for the rational parts of degree 4 polynomials, we define:

$$I_4(\epsilon_1,\epsilon_2,\epsilon_3,\epsilon_4)\equiv\int\frac{\mathrm{d}^D p}{i\pi^{D/2}}\,\frac{(\epsilon_1,p)\,(\epsilon_2,p-k_1)\,(\epsilon_3,p-K_{12})\,(\epsilon_4,p+K_4)}{p^2(p-k_1)^2(p-K_{12})^2(p+K_4)^2},\tag{14}$$

where $K_{12}=k_1+K_2$. For the two-mass-easy case, we have

$$\begin{aligned}I_4(\lambda_3\tilde\lambda_1,\epsilon_2,\lambda_1\tilde\lambda_3,\epsilon_4) &= -\frac{1}{4}\left(\frac{K_2^2+s}{K_2^2-s}+\frac{K_4^2+t}{K_4^2-t}\right)(\epsilon_2,k_1)(\epsilon_4,k_1)\\ &\quad-\frac{1}{4}\left(\frac{K_2^2+t}{K_2^2-t}+\frac{K_4^2+s}{K_4^2-s}\right)(\epsilon_2,k_3)(\epsilon_4,k_3)-\frac{5}{9}\,(k_1,k_3)(\epsilon_2,\epsilon_4)\\ &\quad+\frac{4}{9}\Big((\epsilon_2,k_1)(\epsilon_4,k_3)+(\epsilon_2,k_3)(\epsilon_4,k_1)\Big),\end{aligned}\tag{15}$$

$$\begin{aligned}I_4(\lambda_1\tilde\lambda_3,\epsilon_2,\lambda_1\tilde\lambda_3,\epsilon_4) &= \frac{5}{9}\,\langle 1|\epsilon_2|3]\,\langle 1|\epsilon_4|3]\\ &\quad+\frac{\langle 1|K_2|3]^2}{3}\left[\frac{(\epsilon_2,k_1)\,(\epsilon_4,k_1)}{(K_2^2-s)\,(K_4^2-t)}+\frac{(\epsilon_2,k_3)\,(\epsilon_4,k_3)}{(K_2^2-t)\,(K_4^2-s)}\right].\end{aligned}\tag{16}$$

Other cases can be either obtained by conjugation or by relabelling $k_{1,3}$.

The 3-mass triangle and 2-mass-hard box integrals are not needed for the computation of the 5-gluon amplitudes. The rational parts for these integrals can be found in [1, 32].

5 MHV: $R(1^-2^-3^+4^+5^+)$

Now we begin the computation of the rational parts of the 5-gluon amplitudes. The reader should consult eq. (31) in [1] for the detailed definition of the rational part. In this section we compute the rational part for the helicity configuration $(1^-2^-3^+4^+5^+)$. For this helicity configuration we choose the following polarization vectors:

$$\epsilon_1=\frac{\lambda_1\tilde\lambda_2}{[1\,2]},\qquad \epsilon_2=\frac{\lambda_2\tilde\lambda_1}{[2\,1]},\tag{17}$$

$$\epsilon_3=\frac{\lambda_4\tilde\lambda_3}{\langle 4\,3\rangle},\qquad \epsilon_5=\frac{\lambda_4\tilde\lambda_5}{\langle 4\,5\rangle},\qquad \epsilon_4=\frac{\eta\tilde\lambda_4}{\langle \eta\,4\rangle}.\tag{18}$$

In order to keep the symmetry under $1 \leftrightarrow 2$, $3 \leftrightarrow 5$ manifest, we leave the reference momentum for ϵ_4 arbitrary, i.e. η is an arbitrary (holomorphic) spinor. The final result should be independent of η. In the following formulas for R_i's, we will omit all the denominators of the polarization vectors.

We classify all the 21 diagrams into 7 sets. They are shown in Figs. 5 to 11. Now let us present the results of the rational parts for these 7 sets of Feynman diagrams.



Figure 5: The first diagram is the only 5-point diagram. Its combination with the pinched $k_{1,2} \to k_{12}$ 4-point diagram leads to triangle diagrams only by making use of the reduction formula (5).

For the two Feynman diagrams given in Fig. 5, by using the tensor reduction formula given in eq. (5), they are reduced to just triangle diagrams. Because the other three external lines have the same helicity, these triangle diagrams can be further reduced to bubble diagrams which can be computed easily. We note that some extra terms must be added for tensor reduction of the box or triangle integrals [1]. The final result for the rational part is exceptionally simple and is given as follows:

$$R_1 = -\frac{1}{18}\left((\epsilon_4, \epsilon_5)\epsilon_3 + (\epsilon_3, \epsilon_4)\epsilon_5, k_1 - k_2\right). \tag{19}$$

For the four Feynman diagrams shown in Fig. (6), we can use the same tensor reduction formula as above. Because we use the tensor reduction formula for the box integral, we should add an extra term. The rational part of these four diagrams is:

$$\begin{aligned} R_2 &= \frac{1}{18}\left((\epsilon_{34}, k_2)(\epsilon_5, k_1) + (\epsilon_3, k_2)(\epsilon_{45}, k_1)\right) \\ &+ \frac{1}{18}\left(2s_{51} - s_{34} - 3s_{12}\right)(\epsilon_{34}, \epsilon_5) \end{aligned}$$



Figure 6: These 4 Feynman diagrams are reduced as the 2 diagrams in Fig. 5.

$$+\quad \frac{1}{18}\,(2s_{23}-s_{45}-3s_{12})\,(\epsilon_3,\epsilon_{45}). \tag{20}$$

The four diagrams shown in Fig. 7 can also be reduced simply. Because we leave the reference momentum of ϵ_4 arbitrary, the reduction is just to triangle integrals by using eq. (7). The rational part is:

$$\begin{aligned} R_3 \;=\;& -\frac{1}{9}\,s_{23}(\tilde{\epsilon}_{23},\epsilon_{51}) - \frac{\langle \eta\,4\rangle}{18\,\langle 3\,4\rangle}\,s_{51}(\epsilon_2,\epsilon_{51}) - \frac{1}{6}\,(\eta\tilde{\lambda}_3,k_4)\,(\epsilon_2,\epsilon_{51}) \\ &+\frac{1}{9}\,s_{51}(\tilde{\epsilon}_{51},\epsilon_{23}) - \frac{\langle \eta\,4\rangle}{18\,\langle 5\,4\rangle}\,s_{23}(\epsilon_1,\epsilon_{23}) - \frac{1}{6}\,(\eta\tilde{\lambda}_5,k_4)\,(\epsilon_1,\epsilon_{23}), \end{aligned} \tag{21}$$

where $\tilde{\epsilon}_{23} = \epsilon_{23}|_{\epsilon_3\to\eta\tilde{\lambda}_3}$ and $\tilde{\epsilon}_{51} = \epsilon_{51}|_{\epsilon_5\to\eta\tilde{\lambda}_5}$, i.e.:

$$\tilde{\epsilon}_{23} \;=\; \frac{[1\,3]}{[2\,3]}\,\eta\tilde{\lambda}_3 - \frac{\langle 2\,\eta\rangle}{\langle 2\,3\rangle}\,\lambda_2\tilde{\lambda}_1 + \frac{\langle 2\,\eta\rangle\,[3\,1]}{2\,\langle 2\,3\rangle\,[3\,2]}\,(k_2-k_3), \tag{22}$$

$$\tilde{\epsilon}_{51} \;=\; \frac{\langle 1\,\eta\rangle}{\langle 1\,5\rangle}\,\lambda_1\tilde{\lambda}_2 - \frac{[2\,5]}{[1\,5]}\,\eta\tilde{\lambda}_5 + \frac{\langle 1\,\eta\rangle\,[5\,2]}{2\,\langle 1\,5\rangle\,[5\,1]}\,(k_5-k_1). \tag{23}$$

The Feynman diagram shown in Fig. 8 is the only 2-mass triangle diagram which does not combine with a higher point diagram. This actually gives rise to spurious pole terms which can only be cancelled by the contribution from



(a) (b) (c) (d)

Figure 7: The 3rd set of Feynman diagrams. They can also be reduced easily because of the same adjacent helicity 3^+4^+ or 4^+5^+.



Figure 8: The only 2-mass triangle diagram which does not combine with a higher point diagram.

cut-constructible part. The rational part is computed by using eq. (11):

$$\begin{aligned}
R_4 &= \frac{1}{36}\left(7(\epsilon_4,\epsilon_{51})(\epsilon_{23},k_4) - 7(\epsilon_4,\epsilon_{23})(\epsilon_{51},k_4) + 4(\epsilon_{23},\epsilon_{51})(\epsilon_4,k_{51})\right)\\
&- \frac{s_{51}+s_{23}}{12(s_{51}-s_{23})}\left((\epsilon_4,\epsilon_{51})(\epsilon_{23},k_4) + (\epsilon_4,\epsilon_{23})(\epsilon_{51},k_4)\right)\\
&- \frac{s_{51}+s_{23}}{6(s_{51}-s_{23})^2}(\epsilon_4,k_{51})(\epsilon_{51},k_4)(\epsilon_{23},k_4)\\
&- \frac{1}{4}(\epsilon_4,(\epsilon_5,\epsilon_1)\epsilon_{23} + (\epsilon_2,\epsilon_3)\epsilon_{51})\\
&- \frac{1}{4}\,\frac{(\epsilon_4,k_{51})\left((\epsilon_5,\epsilon_1)\epsilon_{23} + (\epsilon_2,\epsilon_3)\epsilon_{51},k_4\right)}{s_{51}-s_{23}}.
\end{aligned} \tag{24}$$



Figure 9: This set has 2 Feynman diagrams. Their reduction leads to tadpole (one-point) diagrams which are zero in dimensional regularization.

The 2 Feynman diagrams shown in Fig. 9 are reduced to tadpole (one-point) diagrams which are zero in dimensional regularization. The only contribution is from the extra terms and the result is:

$$R_5 \;=\; -\frac{1}{6}\, s_{12}(\epsilon_{345}, k_1) + \frac{1}{4}\, s_{12}\left((\epsilon_3, \epsilon_{45}) + (\epsilon_{34}, \epsilon_5)\right). \tag{25}$$



Figure 10: This set has 4 Feynman diagrams. Tensor reduction by making use of eq. (7) leads to bubble diagrams.

The 4 Feynman diagrams shown in Fig. 10 are reduced to bubble diagrams by making use of eq. (7) because the 2 massless external lines $k_{3,4}$ or $k_{4,5}$ have the same adjacent helicity. The rational part is:

$$\begin{aligned} R_6 \;=\;& (\eta\tilde{\lambda}_3, k_4)\left[\frac{1}{4}\left((\epsilon_5, \epsilon_{12}) + (\epsilon_{51}, \epsilon_2)\right) + \frac{1}{6}\,(\epsilon_{512}, k_4)\right] \\ +\;& (\eta\tilde{\lambda}_5, k_4)\left[\frac{1}{4}\left((\epsilon_1, \epsilon_{23}) + (\epsilon_{12}, \epsilon_3)\right) - \frac{1}{6}\,(\epsilon_{123}, k_4)\right]. \end{aligned} \tag{26}$$

The last 4 diagrams shown in Fig. 11 are two pairs of Feynman diagrams with different adjacent helicities. By explicit computation we found that



Figure 11: The last set has two pairs of Feynman diagrams with different adjacent helicities. Each combination is identically zero.

each combination is identically zero. This result is true for more general cases where the composite polarization vector can be arbitrary.

Having computed all the 21 diagrams separately, the complete rational part is obtained by adding them together. We have

$$
\begin{aligned}
R \;=\;& \frac{1}{[1\,2]^2\,\langle 3\,4\rangle\,\langle 4\,5\rangle\,\langle \eta\,4\rangle}\left[-\frac{1}{18}\left((\epsilon_4,\epsilon_5)\epsilon_3+(\epsilon_3,\epsilon_4)\epsilon_5,k_1-k_2\right)\right.\\
&+\frac{1}{18}\left((\epsilon_{34},k_2)(\epsilon_5,k_1)+(\epsilon_3,k_2)(\epsilon_{45},k_1)\right)\\
&+\frac{1}{18}\left(2s_{51}-s_{34}-3s_{12}\right)(\epsilon_{34},\epsilon_5)+\frac{1}{18}\left(2s_{23}-s_{45}-3s_{12}\right)(\epsilon_3,\epsilon_{45})\\
&-\frac{1}{9}\,s_{23}(\tilde{\epsilon}_{23},\epsilon_{51})-\frac{\langle \eta\,4\rangle}{18\,\langle 3\,4\rangle}\,s_{51}(\epsilon_2,\epsilon_{51})-\frac{1}{6}\,(\eta\tilde{\lambda}_3,k_4)\,(\epsilon_2,\epsilon_{51})\\
&+\frac{1}{9}\,s_{51}(\tilde{\epsilon}_{51},\epsilon_{23})-\frac{\langle \eta\,4\rangle}{18\,\langle 5\,4\rangle}\,s_{23}(\epsilon_1,\epsilon_{23})-\frac{1}{6}\,(\eta\tilde{\lambda}_5,k_4)\,(\epsilon_1,\epsilon_{23})\\
&-\frac{1}{6}\,s_{12}(\epsilon_{345},k_1)+\frac{1}{4}\,s_{12}\left((\epsilon_3,\epsilon_{45})+(\epsilon_{34},\epsilon_5)\right)\\
&+(\eta\tilde{\lambda}_3,k_4)\left[\frac{1}{4}\left((\epsilon_5,\epsilon_{12})+(\epsilon_{51},\epsilon_2)\right)+\frac{1}{6}\,(\epsilon_{512},k_4)\right]\\
&+(\eta\tilde{\lambda}_5,k_4)\left[\frac{1}{4}\left((\epsilon_1,\epsilon_{23})+(\epsilon_{12},\epsilon_3)\right)-\frac{1}{6}\,(\epsilon_{123},k_4)\right]\\
&+\frac{1}{36}\left(7(\epsilon_4,\epsilon_{51})(\epsilon_{23},k_4)-7(\epsilon_4,\epsilon_{23})(\epsilon_{51},k_4)+4(\epsilon_{23},\epsilon_{51})(\epsilon_4,k_{51})\right)\\
&-\frac{s_{51}+s_{23}}{12(s_{51}-s_{23})}\left((\epsilon_4,\epsilon_{51})(\epsilon_{23},k_4)+(\epsilon_4,\epsilon_{23})(\epsilon_{51},k_4)\right)
\end{aligned}
$$

$$
\begin{aligned}
- &\quad \frac{s_{51}+s_{23}}{6(s_{51}-s_{23})^2}\,(\epsilon_4,k_{51})(\epsilon_{51},k_4)(\epsilon_{23},k_4) \\
- &\quad \frac{1}{4}(\epsilon_4,(\epsilon_5,\epsilon_1)\epsilon_{23}+(\epsilon_2,\epsilon_3)\epsilon_{51}) \\
- &\quad \frac{1}{4}\,\frac{(\epsilon_4,k_{51})\,((\epsilon_5,\epsilon_1)\epsilon_{23}+(\epsilon_2,\epsilon_3)\epsilon_{51},k_4)}{s_{51}-s_{23}}\Bigg]\,. \qquad (27)
\end{aligned}
$$

For the same helicity configuration, the rational part of the amplitude obtained by Bern, Dixon and Kosower as given in [2] is :

$$
\begin{aligned}
\tilde{R} &= \frac{2}{9}\,\frac{\langle 1\,2\rangle^4}{\langle 1\,2\rangle\,\langle 2\,3\rangle\,\langle 3\,4\rangle\,\langle 4\,5\rangle\,\langle 5\,1\rangle}-\frac{1}{3}\,\frac{\langle 3\,5\rangle\,[3\,5]^3}{[1\,2]\,[2\,3]\,\langle 3\,4\rangle\,\langle 4\,5\rangle\,[5\,1]} \\
&+ \frac{1}{3}\,\frac{\langle 1\,2\rangle\,[3\,5]^2}{[2\,3]\,\langle 3\,4\rangle\,\langle 4\,5\rangle\,[5\,1]}+\frac{1}{6}\,\frac{\langle 1\,2\rangle\,[3\,4]\,\langle 4\,1\rangle\,\langle 2\,4\rangle\,[4\,5]}{s_{23}\,\langle 3\,4\rangle\,\langle 4\,5\rangle\,s_{51}} \\
&- \frac{1}{6}\,\frac{[3\,4]\,\langle 4\,1\rangle\,\langle 2\,4\rangle\,[4\,5]\,(\langle 2\,3\rangle\,[3\,4]\,\langle 4\,1\rangle+\langle 2\,4\rangle\,[4\,5]\,\langle 5\,1\rangle)}{s_{23}\,\langle 3\,4\rangle\,\langle 4\,5\rangle\,s_{51}} \\
&\quad \times\frac{s_{51}+s_{23}}{(s_{51}-s_{23})^2}\,, \qquad (28)
\end{aligned}
$$

which can be obtained from the full amplitude eq. (8) and (10) in [2] by making the replacement

$$
\begin{aligned}
\mathrm{L}_0(\mathrm{r}) &\to 0\,, \\
\mathrm{L}_1(\mathrm{r}) &\to \frac{1}{1-r}\,, \\
\mathrm{L}_2(\mathrm{r}) &\to \frac{1}{2}\frac{(r+1)}{r(1-r)^2} \qquad (29)
\end{aligned}
$$

and discarding V^f which is absorbed in the cut constructible part. One can find detailed discussions about how to construct the rational part from Bern etc.'s results in [28, 30]. We did not find an easy way to prove that these results (R and $\tilde{R}$) agree with each other. With the help of Mathematica one can easily check the following result:

$$
R = \frac{1}{2}\,\tilde{R}. \qquad (30)
$$

This shows that the rational part computed directly from Feynman integrals agrees with the well-known result of Bern, Dixon and Kosower [2]. The factor

1/2 is because R is for the amplitude with a real scalar circulating in the loop according to our definition in eq. (31) in [1], while in their notation $\tilde{R}$ is for the amplitude with a complex scalar.

6 MHV: $R(1^-2^+3^-4^+5^+)$

For this helicity configuration we choose the following polarization vectors:

$$\epsilon_1 = \frac{\lambda_1\tilde{\lambda}_5}{[1\,5]}, \qquad \epsilon_3 = \frac{\lambda_3\tilde{\lambda}_4}{[3\,4]}, \tag{31}$$

$$\epsilon_4 = \frac{\lambda_5\tilde{\lambda}_4}{\langle 5\,4\rangle}, \qquad \epsilon_5 = \frac{\lambda_4\tilde{\lambda}_5}{\langle 4\,5\rangle}, \qquad \epsilon_2 = \frac{\eta\tilde{\lambda}_2}{\langle \eta\,2\rangle}. \tag{32}$$

The reference momentum of ϵ_2 is arbitrary. As before, in the following formulas for R_i's, we will omit all the denominators of the polarization vectors.

The 21 Feynman diagrams are classified into 6 sets. Apart from the last set, they are shown in Figs. 12 to 16. Let us compute the rational part from each set in turn.



Figure 12: 6 Feynman diagrams with the same adjacent helicity.

The 1st set consists of 2 Feynman diagrams which include the only pentagon diagram. Tensor reduction is easy by using eq. (5) which gives rise to triangle diagrams. The rational parts of the reduced integrals can be computed by using eqs. (9) to (11) and we have:

$$\begin{aligned} R_1 &= -\frac{1}{18}\left((\epsilon_2,\epsilon_3)\epsilon_1 + (\epsilon_1,\epsilon_2)\epsilon_3 + (\epsilon_1,\epsilon_3)\epsilon_2, k_4 - k_5\right) \\ &+ \frac{1}{12}(\epsilon_1,\epsilon_2)(\epsilon_3, k_2 - k_{45}) - \frac{1}{12}(\epsilon_2,\epsilon_3)(\epsilon_1, k_2 - k_{45}) \end{aligned}$$

$$\begin{aligned} &+ \quad \frac{s_{34}+s_{51}}{12(s_{34}-s_{51})}\,(((\epsilon_2,\epsilon_3)\epsilon_1+(\epsilon_2,\epsilon_1)\epsilon_3,k_2) \\ &+ \quad \frac{s_{34}+s_{51}}{6(s_{34}-s_{51})^2}\,(\epsilon_2,k_{34})(\epsilon_3,k_2)(\epsilon_1,k_2). \end{aligned} \tag{33}$$



Figure 13: This set consists of 4 Feynman diagrams which can be reduced easily because of the same adjacent helicity.

The 2nd set of Feynman diagrams consist of 4 diagrams. They are reduced identically as in the above. We only need to add an extra term because of the box reduction in $D = 4$. The rational part from set 2 is:

$$\begin{aligned} R_2 \quad = \quad & \left(-\frac{1}{6}\,s_{45}+\frac{1}{9}\,s_{51}-\frac{1}{18}\,s_{23}\right)(\epsilon_1,\epsilon_{23}) \\ + \quad & \left(-\frac{1}{6}\,s_{45}+\frac{1}{9}\,s_{34}-\frac{1}{18}\,s_{12}\right)(\epsilon_{12},\epsilon_3) \\ + \quad & \frac{1}{18}\,((\epsilon_{23},k_4)(\epsilon_1,k_5)+(\epsilon_3,k_4)(\epsilon_{12},k_5)). \end{aligned} \tag{34}$$

The two diagrams shown in Fig. 14 are the most complicated for the 5-gluon amplitude. There is no simple reduction formula to simplify the computation. One needs to use the 2-mass-easy formulas as given in Sect. 4. The explicit result of the rational part from the second diagram is:

$$\begin{aligned} R_3(2) \quad = \quad & \frac{\langle\eta\,4\rangle\,\langle 5\,2\rangle}{18\,\langle 2\,4\rangle\,\langle 4\,2\rangle}\,[(\epsilon_3,k_2)(\epsilon_{51},k_4)-2(k_2,k_4)(\epsilon_3,\epsilon_{51})] \\ + \quad & \frac{\langle\eta\,4\rangle}{\langle 2\,4\rangle}\left[\frac{1}{9}\,((\epsilon_4,\epsilon_{51})\epsilon_3+(\epsilon_3,\epsilon_{51})\epsilon_4,k_2)+\frac{(\epsilon_4,k_{51})\,(\epsilon_{51},k_4)\,(\epsilon_3,k_2)}{6\,(s_{51}-s_{23})}\right] \end{aligned}$$



Figure 14: 2 Feynman diagrams which are computed by using the 2-mass-easy formulas as given in Sect. 4.

$$
\begin{aligned}
+ \quad & \frac{\langle 5\,2\rangle}{\langle 4\,2\rangle}\left[\frac{1}{12}\left((\epsilon_2,\epsilon_3)\,\epsilon_{51}-(\epsilon_2,\epsilon_{51})\,\epsilon_3,k_2\right)\right. \\
& +\frac{1}{18}(2(\epsilon_3,\epsilon_{51})\epsilon_2-(\epsilon_2,\epsilon_3)\epsilon_{51},k_4) \\
& -\frac{s_{34}+s_{51}}{12\,(s_{34}-s_{51})}\left((\epsilon_2,\epsilon_3)\,\epsilon_{51}+(\epsilon_2,\epsilon_{51})\,\epsilon_3,k_2\right) \\
& \left.+\frac{(s_{34}+s_{51})}{6\,(s_{34}-s_{51})^2}\,(\epsilon_2,k_{51})\,(\epsilon_3,k_2)\,(\epsilon_{51},k_2)\right] \\
+ \quad & \frac{\langle \eta\,2\rangle\,\langle 5\,4\rangle}{\langle 4\,2\rangle\,\langle 2\,4\rangle}\left[-\frac{5}{9}(\epsilon_3,\epsilon_{51})(k_2,k_4)+\frac{4}{9}(\epsilon_3,k_2)(\epsilon_{51},k_4)\right. \\
& \left.+\frac{1}{4}(\epsilon_3,k_2)(\epsilon_{51},k_2)\left(1+\frac{s_{34}+s_{51}}{s_{34}-s_{51}}\right)\right]. \qquad (35)
\end{aligned}
$$

The rational part from the first diagram in Fig. 14 can be obtained from the above result by the symmetry transformation:

$$R_3(1) = -R_3(2)|_{1\leftrightarrow 3, 4\leftrightarrow 5}. \qquad (36)$$

We also set $R_3 = R_3(1) + R_3(2)$.



Figure 15: The 4th set consists of 3 two-mass triangle diagrams.

The rational part of the 3 two-mass triangle diagrams shown in Fig. 15 can be directly computed by using eqs. (11) and (10). The rational part from the first diagram in Fig. 15 is denoted by $R_4(0)$ and is given by:

$$\begin{aligned} R_4(0) &= \frac{1}{36}\left(7(\epsilon_2,\epsilon_{34})(\epsilon_{51},k_2) - 7(\epsilon_2,\epsilon_{51})(\epsilon_{34},k_2) + 4(\epsilon_{34},\epsilon_{51})(\epsilon_2,k_{34})\right) \\ &- \frac{s_{34}+s_{51}}{12(s_{34}-s_{51})}\left((\epsilon_2,\epsilon_{34})(\epsilon_{51},k_2) + (\epsilon_2,\epsilon_{51})(\epsilon_{34},k_2)\right) \\ &- \frac{s_{34}+s_{51}}{6(s_{34}-s_{51})^2}(\epsilon_2,k_{34})(\epsilon_{34},k_2)(\epsilon_{51},k_2). \end{aligned} \tag{37}$$

The rational part from the third Feynman diagram in Fig. 15 is denoted by $R_4(1)$ and we have:

$$\begin{aligned} R_4(1) &= \frac{1}{36}\left(7(\epsilon_5,\epsilon_{12})(\epsilon_{34},k_5) - 7(\epsilon_5,\epsilon_{34})(\epsilon_{12},k_5) + 4(\epsilon_{12},\epsilon_{34})(\epsilon_5,k_{12})\right) \\ &- \frac{s_{12}+s_{34}}{12(s_{12}-s_{34})}\left((\epsilon_5,\epsilon_{12})(\epsilon_{34},k_5) + (\epsilon_5,\epsilon_{34})(\epsilon_{12},k_5)\right) \\ &- \frac{s_{12}+s_{34}}{6(s_{12}-s_{34})^2}(\epsilon_5,k_{12})(\epsilon_{12},k_5)(\epsilon_{34},k_5) \\ &- \frac{1}{4}(\epsilon_1,\epsilon_2)\left[(\epsilon_5,\epsilon_{34}) + \frac{(\epsilon_5,k_{12})(\epsilon_{34},k_5)}{s_{12}-s_{34}}\right]. \end{aligned} \tag{38}$$

The rational part from the second diagram in Fig. 15 is denoted by $R_4(2)$ and it can be obtained from $R_4(1)$ by symmetry operation. In total we have:

$$R_4 = R_4(0) + R_4(1) - (R_4(1)|_{1\leftrightarrow 3, 4\leftrightarrow 5}). \tag{39}$$



Figure 16: 2 Feynman diagrams with the same adjacent helicity $k_{4,5}$.

The 5th set consists of 2 Feynman diagrams which are reduced to tadpoles. The rational part is:

$$R_5 = -\frac{1}{6}\,(\epsilon_{123}, k_4)\, s_{45} + \frac{1}{4}\,((\epsilon_1, \epsilon_{23}) + (\epsilon_{12}, \epsilon_3))\, s_{45}. \tag{40}$$

Up to now we have computed 13 Feynman diagrams. The remaining 8 Feynman diagrams consist of 4 one-mass triangle diagrams and 4 bubble diagrams. As we showed in last section, they are identically zero for each pair of triangle and bubble diagrams.

The final result for the rational part is:

$$R \;=\; -\frac{1}{[1\,5]\,[3\,4]\,\langle 4\,5\rangle^2\,\langle \eta\,2\rangle}\sum_{i=1}^{5} R_i. \tag{41}$$

In comparison, the result obtained by Bern, Dixon and Kosower [2] from string theory is:

$$\begin{aligned}
\tilde{R} \;=\;& \frac{2}{9}\,\frac{\langle 1\,3\rangle^4}{\langle 1\,2\rangle\,\langle 2\,3\rangle\,\langle 3\,4\rangle\,\langle 4\,5\rangle\,\langle 5\,1\rangle} + \frac{1}{3}\,\frac{[2\,4]^2\,[2\,5]^2}{[1\,2]\,[2\,3]\,[3\,4]\,\langle 4\,5\rangle\,[5\,1]} \\
-\;& \frac{1}{3}\,\frac{\langle 1\,2\rangle\,\langle 4\,1\rangle^2\,[2\,4]^3}{\langle 4\,5\rangle\,\langle 5\,1\rangle\,\langle 2\,4\rangle\,[2\,3]\,[3\,4]\,s_{51}} + \frac{1}{3}\,\frac{\langle 3\,2\rangle\,\langle 5\,3\rangle^2\,[2\,5]^3}{\langle 5\,4\rangle\,\langle 4\,3\rangle\,\langle 2\,5\rangle\,[2\,1]\,[1\,5]\,s_{34}} \\
+\;& \frac{1}{6}\,\frac{\langle 1\,3\rangle^2\,[2\,4]\,[2\,5]}{s_{34}\,\langle 4\,5\rangle\,s_{51}} - \frac{\langle 1\,2\rangle\,\langle 2\,3\rangle\,\langle 3\,4\rangle\,\langle 4\,1\rangle^2\,[2\,4]^2}{\langle 4\,5\rangle\,\langle 5\,1\rangle\,\langle 2\,4\rangle^2\,s_{51}}\left[\frac{1}{s_{51}-s_{23}} - \frac{1}{s_{34}-s_{51}}\right] \\
+\;& \frac{\langle 3\,2\rangle\,\langle 2\,1\rangle\,\langle 1\,5\rangle\,\langle 5\,3\rangle^2\,[2\,5]^2}{\langle 5\,4\rangle\,\langle 4\,3\rangle\,\langle 2\,5\rangle^2\,s_{34}}\left[\frac{1}{s_{34}-s_{51}} - \frac{1}{s_{12}-s_{34}}\right] \\
+\;& \frac{1}{3}\,\frac{\langle 2\,3\rangle^2\,\langle 4\,1\rangle^3\,[2\,4]^3}{\langle 4\,5\rangle\,\langle 5\,1\rangle\,\langle 2\,4\rangle\,s_{23}\,s_{51}}\,\frac{s_{51}+s_{23}}{(s_{51}-s_{23})^2} \\
-\;& \frac{1}{3}\,\frac{\langle 2\,1\rangle^2\,\langle 5\,3\rangle^3\,[2\,5]^3}{\langle 5\,4\rangle\,\langle 4\,3\rangle\,\langle 2\,5\rangle\,s_{12}\,s_{34}}\,\frac{s_{12}+s_{34}}{(s_{12}-s_{34})^2} \\
+\;& \left[\frac{1}{6}\,\frac{\langle 1\,3\rangle\,[2\,4]\,[2\,5]\,(\langle 1\,5\rangle\,[5\,2]\,\langle 2\,3\rangle - \langle 3\,4\rangle\,[4\,2]\,\langle 2\,1\rangle)}{\langle 4\,5\rangle}\right. \\
& \left. + \frac{1}{3}\,\frac{\langle 1\,2\rangle^2\,\langle 3\,4\rangle^2\,\langle 4\,1\rangle\,[2\,4]^3}{\langle 4\,5\rangle\,\langle 5\,1\rangle\,\langle 2\,4\rangle} - \frac{1}{3}\,\frac{\langle 3\,2\rangle^2\,\langle 1\,5\rangle^2\,\langle 5\,3\rangle\,[2\,5]^3}{\langle 5\,4\rangle\,\langle 4\,3\rangle\,\langle 2\,5\rangle}\right] \\
& \times \frac{s_{34}+s_{51}}{s_{34}\,s_{51}\,(s_{34}-s_{51})^2}.
\end{aligned} \tag{42}$$

As in the other MHV case, we did not find an easy way to prove that these results (R and $\tilde{R}$) agree with each other. With the help of Mathematica one can easily check the following result:

$$R = \frac{1}{2}\,\tilde{R}. \tag{43}$$

This shows that the rational part computed directly from Feynman integrals agrees with the well-known result of Bern, Dixon and Kosower [2].

7 Conclusion

In this paper, by using the formalism developed in the previous paper [1], we computed explicitly the rational parts of the 5-gluon one-loop amplitudes for the 2 MHV helicity configurations. The results are in agreement with the well-known results of Bern, Dixon and Kosower [2]. Intermediate results for some combinations of Feynman diagrams are presented explicitly. The explicit result for the MHV configuration $(- - + + +)$ as given in eq. (27) is more complicated than the previous known result eq. (28). In fact the use of the composite polarization vectors hides some more complexities of our results. The correctness and usefulness of the formalism as developed in [1] will be further strengthed in [32] by computing the rational parts of the 6-gluon one-loop amplitudes for all possible helicity configurations.

Acknowledgments

CJZ would like to thank J. -P. Ma for constant encouragements, helpful discussions and careful reading of the paper. His (financial) support (to buy a computer which was still in use today) actually goes back to the much earlier difficult times when I did not have enough grants from other sources. What is more important is that there are no strings attached to his support and it is up to the last author to explore what he wants to. CJZ would also like to thank R. Iengo for encouragements and his interests in this work, helpful discussions and comments; to C. F. Berger, Z. Bern, Z. Chang, L. J. Dixon, B. Feng, D. Forde, E. Gava, H. -Y. Guo, D. A. Kosower, K. S. Narain, K. Wu, Y. -S. Wu, Z. Xu and Z. -X. Zhang for discussions and comments; to Prof. X. -Q. Li and the hospitality at Nankai University where we can have good food; and finally to Prof. S. Randjbar-Daemi and the hospitality at Abdus Salam International Center for Theoretical Physics, Trieste, Italy. This work is supported in part by funds from the National Natural Science Foundation of China with grant number 10475104 and 10525522.

References

[1] Z. -G. Xiao, G. Yang and C. -J. Zhu, "The rational parts of one-loop QCD amplitudes I: The general formalism," hep-ph/0607015.

[2] Z. Bern, L. J. Dixon and D. A. Kosower, "One Loop Corrections to Five Gluon Amplitudes," Phys. Rev. Lett. **70** (1993) 2677-2680, hep-ph/9302280;

Z. Bern, "String based perturbative methods for gauge theories," TASI 92 lectures, hep-ph/9304249.

[3] Z. Kunszt, A. Signer and Z. Trócsányi, "One-loop radiative corrections to the helicity amplitudes of QCD processes involving four quarks and one gluon," Phys. Lett. **B336** (1994) 529-536, hep-ph/9405386.

[4] Z. Bern, L. J. Dixon and D. A. Kosower, "One-loop corrections to two-quark three-gluon amplitudes," Nucl. Phys. **B437** (1995) 259-304, hep-ph/9409393.

[5] R. K. Ellis and J, C. Sexton, "QCD Radiative Corrections to Parton-Parton Scattering," Nucl. Phys. **B269** (1986) 445-484.

[6] Z. Bern and D. A. Kosower, "Efficient Computation Of Gauge Theory Loops Using String Theory," in Copenhagen 1987, Proceedings, Perspectives In String Theory, p. 390-402; "A New Approach To One Loop Calculations In Gauge Theories," Phys. Rev. **D38** (1988) 1888; "Efficient calculation of one loop QCD amplitudes," Phys. Rev. Lett. **66** (1991) 1669-1672; "The Computation of loop amplitudes in gauge theories," Nucl. Phys. **B379** (1992) 451-561.

[7] G. P. Salam, "Developments in perturbative QCD," Invited talk at 22nd International Symposium on Lepton-Photon Interactions at High

Energy (LP 2005), Uppsala, Sweden, 30 Jun - 5 Jul 2005, hep-ph/0510090.

[8] E. Witten, "Perturbative gauge theory as a string theory in twistor space," Commun. Math. Phys. **252** (2004) 189-258, hep-th/0312171.

[9] F. Cachazo, P. Svrček and E. Witten, "MHV vertices and tree amplitudes in gauge theory," JHEP **0409**, (2004) 006, hep-th/0403047.

[10] Z. Bern, L. Dixon, D. C. Dunbar and D. A. Kosower, "One-Loop n-Point Gauge Theory Amplitudes, Unitarity and Collinear Limits," Nucl. Phys. **B425** (1994) 217-260, hep-ph/9403226; "Fusing Gauge Theory Tree Amplitudes Into Loop Amplitudes," Nucl. Phys. **B435** (1995) 59-101, hep-ph/9409265.

[11] A. Brandhuber, B. Spence and G. Travaglini, "One-Loop Gauge Theory Amplitudes In N = 4 Super Yang-Mills From MHV Vertices," Nucl. Phys. B **706** (2005) 150-180, hep-th/0407214; "From Trees to Loops and Back," JHEP **0601** (2006) 142, hep-th/0510253.

[12] F. Cachazo, "Holomorphic Anomaly Of Unitarity Cuts And One-Loop Gauge Theory Amplitudes," hep-th/0410077.

[13] Z. Bern, V. Del Duca, L. J. Dixon and D. A. Kosower, "All Non-Maximally-Helicity-Violating One-Loop Seven-Gluon Amplitudes in N=4 Super-Yang-Mills Theory," Phys. Rev. **D71** (2005) 045006, hep-th/0410224;

S. J. Bidder, N. E. J. Bjerrum-Bohr, L, J. Dixon and D. C. Dunbar, "N=1 Supersymmetric One-loop Amplitudes and the Holomorphic Anomaly of Unitarity Cuts," Phys. Lett. **B606** (2005) 189-201, hep-th/0410296;

Z. Bern, L. J. Dixon and D. A. Kosower, "All Next-to-Maximally-Helicity-Violating One-Loop Gluon Amplitudes in N=4 Super-Yang-Mills Theory," Phys. Rev. **D72** (2005) 045014, hep-th/0412210.

[14] C. Quigley and M. Rozali, "One-Loop MHV Amplitudes in Supersymmetric Gauge Theories," JHEP **0501** (2005) 053, hep-th/0410278.

[15] J. Bedford, A. Brandhuber, B. Spence and G. Travaglini, “A Twistor Approach to One-Loop Amplitudes in N=1 Supersymmetric Yang-Mills Theory,” Nucl. Phys. **B706** (2005) 100-126, hep-th/0410280; “Non-Supersymmetric Loop Amplitudes and MHV Vertices,” Nucl. Phys. **B712** (2005) 59-85, hep-th/0412108.

[16] S. J. Bidder, N.E.J. Bjerrum-Bohr, D. C. Dunbar and W. B. Perkins, “Twistor Space Structure of the Box Coefficients of N=1 One-loop Amplitudes,” Phys. Lett. **B608** (2005) 151-163, hep-th/0412023;

S. J. Bidder, N. E. J. Bjerrum-Bohr, D. C. Dunbar and W. B. Perkins, “One-Loop Gluon Scattering Amplitudes in Theories with $N < 4$ Supersymmetries,” Phys. Lett. **B612** (2005) 75-88, hep-th/0502028.

[17] Z. Bern, N. E. J. Bjerrum-Bohr, D. C. Dunbar and H. Ita, “Recursive Calculation of One-Loop QCD Integral Coefficients,” JHEP **0511** (2005) 027, hep-ph/0507019.

[18] M. -X. Luo and C. -K. Wen, “One-Loop Maximal Helicity Violating Amplitudes in N=4 Super Yang-Mills Theories,” JHEP **0411** (2004) 004, hep-th/0410045; “Systematics of One-Loop Scattering Amplitudes in N=4 Super Yang-Mills Theories,” Phys. Lett. **B609** (2005) 86-94, hep-th/0410118.

[19] R. Britto, F. Cachazo and B. Feng, “New Recursion Relations for Tree Amplitudes of Gluons,” Nucl. Phys. **B715** (2005) 499-522, hep-th/0412308;

R. Britto, F. Cachazo, B. Feng and E. Witten, “Direct Proof Of Tree-Level Recursion Relation In Yang-Mills Theory,” Phys. Rev. Lett. **94** (2005) 181602, hep-th/0501052.

[20] R. Britto, F. Cachazo and B. Feng, “Generalized Unitarity and One-Loop Amplitudes in N=4 Super-Yang-Mills,” Nucl.Phys. **B725** (2005) 275-305, hep-th/0412103.

[21] R. Britto, E. Buchbinder, F. Cachazo, B. Feng, “One-Loop Amplitudes Of Gluons In SQCD,” Phys. Rev. **D72** (2005) 065012, hep-ph/0503132.

[22] R. Britto, B. Feng and P. Mastrolia, “The Cut-Constructible Part of QCD Amplitudes,” Phys. Rev. **D73** (2006) 105004, hep-ph/0602178.

[23] Z. Bern, L. Dixon and D. A. Kosower, “One-loop amplitudes for e^+e^- to four partons,” Nucl. Phys. **B513** (1998) 3–86, hep-ph/9708239.

[24] Z. Bern and A. G. Morgan, “Massive Loop Amplitudes from Unitarity,” Nucl. Phys. **B467** (1996) 479-509, hep-ph/9511336.

[25] For reviews see: Z. Bern, L. Dixon and D. A. Kosower, “Progress in One-Loop QCD Computations,” Ann. Rev. Nucl. Part. Sci. **46** (1996) 109-148, hep-ph/9602280; “Unitarity-based Techniques for One-Loop Calculations in QCD,” Nucl. Phys. Proc. Suppl. **51C** (1996) 243-249, hep-ph/9606378.

[26] A. Brandhuber, S. McNamara, B. Spence and G. Travaglini, “Loop Amplitudes in Pure Yang-Mills from Generalised Unitarity,” JHEP **0510** (2005) 011, hep-th/0506068.

[27] Z. Bern, L. J. Dixon and D. A. Kosower, “On-Shell Recurrence Relations for One-Loop QCD Amplitudes,” Phys. Rev. **D71** (2005) 105013, hep-th/0501240; “The Last of the Finite Loop Amplitudes in QCD,” Phys. Rev. **D72** (2005) 125003, hep-ph/0505055.

[28] Z. Bern, L. J. Dixon and D. A. Kosower, “Bootstrapping Multi-Parton Loop Amplitudes in QCD,” Phys. Rev. **D73** (2006) 065013, hep-ph/0507005.

[29] D. Forde and D. A. Kosower, “All-Multiplicity One-Loop Corrections to MHV Amplitudes in QCD,” Phys. Rev. **D73** (2006) 061701, hep-ph/0509358.

[30] C. F. Berger, Z. Bern, L. J. Dixon, D. Forde and D. A. Kosower, “Bootstrapping One-Loop QCD Amplitudes with General Helicities,” hep-ph/0604195.

[31] C. F. Berger, Z. Bern, L. J. Dixon, D. Forde and D. A. Kosower, “All One-loop Maximally Helicity Violating Gluonic Amplitudes in QCD,” hep-ph/0607014.

[32] Z. -G. Xiao, G. Yang and C. -J. Zhu, “The rational parts of one-loop QCD amplitudes III: The 6-gluon case,” hep-ph/0607017.

[33] A. Berends and W. T. Giele, “Recursive Calculations For Processes With N Gluons,” Nucl. Phys. **B306** (1988) 759.

[34] G. Passarino and M. J. G. Veltman, “One-loop Corrections for e^+e^- annihilation into $\mu^+\mu^-$ in the Weinberg modes,” Nucl. Phys. **B160** (1979) 151-207.

[35] D. B. Melrose, “Reduction Of Feynman Diagrams,” Nuovo Cim. **40**, 181 (1965); W. L. van Neerven and J. A. M. Vermaseren, “Large Loop Integrals,” Phys. Lett. B **137**, 241 (1984).

[36] Z. Bern, L. Dixon and D. A. Kosower, “Dimensionally Regulated One-Loop Integrals,” Phys.Lett. **B302** (1993) 299-308, Erratum-ibid. **B318** (1993) 649, hep-ph/9212308; “Dimensionally Regulated Pentagon Integrals,” Nucl. Phys. **B412** (1994) 751-816, hep-ph/9306240.

[37] T. Binoth, J.Ph. Guillet, G. Heinrich, “Reduction formalism for dimensionally regulated one-loop N-point integrals,” Nucl.Phys. **B572** (2000) 361-386, hep-ph/9911342.

[38] O. V. Tarasov, “Connection between Feynman integrals having different values of the space-time dimension,” Phys. Rev. D **54**, 6479 (1996), hep-th/9606018; J. Fleischer, F. Jegerlehner and O. V. Tarasov, “Algebraic reduction of one-loop Feynman graph amplitudes,” Nucl. Phys. B **566**, 423 (2000), hep-ph/9907327.

[39] G. Duplancic and B. Nizic, “Reduction method for dimensionally regulated one-loop N-point Feynman integrals,” Eur. Phys. J. C **35**, 105 (2004), hep-ph/0303184.

[40] A. Denner and S. Dittmaier, “Reduction of one-loop tensor 5-point integrals,” Nucl. Phys. B **658**, 175 (2003), hep-ph/0212259; “Reduction schemes for one-loop tensor integrals,” Nucl. Phys. B **734**, 62 (2006), hep-ph/0509141 and the references therein.

[41] Z. Xu, D.-H. Zhang and L. Chang, “Helicity Amplitudes For Multiple Bremsstrahlung In Massless Nonabelian Theories,” Nucl. Phys. **B291** (1987) 392.

[42] L. J. Dixon, "Calculating Scattering Amplitudes Efficiently," hep-ph/9601359.